



ELSEVIER

ORIGINAL ARTICLE

JOURNAL OF CARDIOLOGY

Official Journal of the Japanese College of Cardiology

www.elsevier.com/locate/jjcc

Positive remodeling is associated with more plaque vulnerability and higher frequency of plaque prolapse accompanied with post-procedural cardiac enzyme elevation compared with intermediate/negative remodeling in patients with acute myocardial infarction

Young Joon Hong (MD), Myung Ho Jeong (MD)*, Yun Ha Choi (RN),
Jum Suk Ko (MD), Min Goo Lee (MD), Won Yu Kang (MD),
Shin Eun Lee (MD), Soo Hyun Kim (MD), Keun Ho Park (MD),
Doo Sun Sim (MD), Nam Sik Yoon (MD), Hyun Ju Youn (MD),
Kye Hun Kim (MD), Hyung Wook Park (MD), Ju Han Kim (MD),
Youngkeun Ahn (MD), Jeong Gwan Cho (MD),
Jong Chun Park (MD), Jung Chae Kang (MD)

Heart Center of Chonnam National University Hospital, Chonnam National University Research Institute of Medical Sciences, Gwangju, South Korea

Received 10 November 2008; received in revised form 27 November 2008; accepted 1 December 2008
Available online 13 February 2009

KEYWORDS

Acute myocardial
infarction;
Remodeling;
Plaque;
Intravascular ultrasound

Summary

Background: We assessed the impact of remodeling patterns on pre- and post-procedural intravascular ultrasound (IVUS) findings and cardiac enzyme elevation after stenting in 310 acute myocardial infarction (AMI) patients.

Methods: The positive remodeling (PR) (PR group, $n = 113$) was defined as remodeling index (lesion/reference external elastic membrane cross-sectional area) > 1.05 , intermediate remodeling (IR) as between 0.95 and 1.05, and negative remodeling (NR) as < 0.95 (IR/NR group, $n = 197$). IVUS findings included ruptured plaque (a cavity that communicated with the lumen with an overlying residual fibrous cap fragment),

* Corresponding author. The Heart Center of Chonnam National University Hospital, 671 Jaebongro, Dong-gu, Gwangju 501-757, South Korea. Tel.: +82 62 220 6243; fax: +82 62 228 7174.

E-mail address: myungho@chollian.net (M.H. Jeong).

multiple ruptured plaques (different plaque ruptures separated by a >5-mm length of artery containing smooth lumen contours), thrombus (discrete intraluminal filling defects), and plaque prolapse (tissue extrusion through the stent strut at post-stenting). We compared pre- and post-procedural IVUS findings and cardiac-specific troponin I (cTnI) elevation after stenting according to the remodeling pattern.

Results: The plaque rupture (60% vs. 42%, $p=0.004$), multiple plaque ruptures (22% vs. 14%, $p=0.014$), and IVUS-detected thrombus (42% vs. 28%, $p=0.012$) were more common in the PR group compared with the IR/NR group. Post-stenting plaque prolapse was observed more frequently (36% vs. 22%, $p=0.008$), and cTnI was elevated more significantly after stenting in the PR group compared with the IR/NR group (Δ cTnI; $+7.8 \pm 51.1$ ng/ml vs. $+0.9 \pm 41.1$ ng/ml, $p=0.008$). Multivariate analysis showed that PR [odds ratio (OR)=1.92; 95% CI 1.04–2.98, $p=0.028$], plaque rupture (OR 1.98; 95% CI 1.16–3.45, $p=0.025$), IVUS-detected thrombus (OR 2.30; 95% CI 1.22–3.98, $p=0.008$), and plaque prolapse (OR 8.40; 95% CI 4.19–16.84, $p<0.001$) were independently associated with post-stenting cTnI elevation.

Conclusions: AMI patients with PR have more plaque vulnerability and higher frequency of plaque prolapse accompanied by post-procedural cardiac enzyme elevation compared with AMI patients with IR/NR.

© 2009 Japanese College of Cardiology. Published by Elsevier Ireland Ltd. All rights reserved.

Introduction

Patterns of arterial remodeling during the course of plaque development have been shown to play a pivotal role in the progression of atherosclerosis [1–5]. In vivo assessment of lesion site remodeling using intravascular ultrasound (IVUS) is performed by comparing the vessel area of the lesion to the reference segment [6]. Arterial remodeling is a heterogeneous process ranging from positive remodeling (PR), which is usually responsible for acute coronary syndromes, to negative remodeling (NR) which is usually associated with stenoses in patients with stable angina [7,8]. Lesions are classified as PR because the lesion site external elastic membrane (EEM) cross-sectional area (CSA) is larger than the reference segment, and lesions are classified as NR because the lesion site EEM CSA is smaller than the reference segment.

Typical IVUS features of acute myocardial infarction (AMI) include PR, plaque rupture, thrombus, and either spotty or deep calcium within the minimum lumen site [9–16]. Several studies have reported that rupture of a vulnerable plaque and subsequent thrombus formation is the most important mechanism leading to an acute coronary syndrome [17,18]. So far, data on the pre-procedural IVUS plaque characteristics and post-procedural IVUS findings according to the remodeling pattern in patients with AMI are lacking. We used IVUS to assess the relation between remodeling pattern and pre- and post-procedural IVUS findings in AMI.

Methods

Study population

From January 9, 2001 to July 31, 2007, we identified a total of 310 patients with a first AMI who underwent pre-intervention IVUS within 24 h from symptom onset, were stented successfully, and had post-intervention IVUS imaging. We excluded patients with prior MI, stent thrombosis, in-stent restenosis lesion, coronary artery bypass graft failure, severe heart failure or cardiogenic shock, patients studied with IVUS more than 24 h after symptom onset, and patients in whom adequate IVUS images could not be obtained. All 310 infarct lesions were treated with stent implantation: 187 with drug-eluting stents [138 with sirolimus-eluting stents (Cypher stent, Cordis, Johnson and Johnson, Miami Lakes, FL, USA) and 49 with paclitaxel-eluting stents (Taxus stent, Boston Scientific, Boston, MA, USA)], and 123 with bare metal stents.

The presence of ST-segment elevation MI was determined by >30 min of continuous chest pain, a new ST-segment elevation ≥ 2 mm on at least two contiguous electrocardiographic leads, and creatine kinase-MB >3 times normal. The presence of non-ST-segment elevation MI was diagnosed by chest pain and a positive cardiac biochemical marker without new ST-segment elevation. Infarct-related arteries were identified using a combination of electrocardiographic findings, left ventricular wall motion abnormalities on left

ventricular angiogram or two-dimensional echocardiogram, and coronary angiogram. The protocol was approved by the institutional review board. Hospital records of patients were reviewed to obtain information on clinical demographics.

Laboratory analysis

Venous blood samples were obtained within 24 h after stenting. The blood samples were centrifuged, and serum was removed and stored at -70°C until the assay could be performed. Cardiac-specific troponin I (cTnI) levels were measured by a paramagnetic particle, chemiluminescent immunoassay (Beckman, Coulter Inc., Fullerton, CA, USA). The serum levels of total cholesterol, triglyceride, low-density lipoprotein cholesterol, and high-density lipoprotein cholesterol were measured by standard enzymatic methods. High-sensitivity C-reactive protein was analyzed turbidimetrically with sheep antibodies against human C-reactive protein; this has been validated against the Dade-Behring method [19].

Quantitative coronary angiography (QCA) analysis

Coronary angiogram was analyzed with validated QCA system (Philips H5000 or Allura DCI program, Philips Medical Systems, Eindhoven, the Netherlands). With the outer diameter of the contrast-filled catheter as the calibration standard, the minimal lumen diameter and reference diameter were measured in diastolic frames from orthogonal projections.

IVUS imaging and analysis

All IVUS examinations were performed before and after stenting after intracoronary administration of $300\text{ }\mu\text{g}$ nitroglycerin using a commercially available IVUS system (Boston Scientific Corporation/SCIMed, Minneapolis, MN, USA). The IVUS catheter was advanced distal to the target lesion, and imaging was performed retrograde to the aorto-ostial junction at an automatic pullback speed of 0.5 mm/sec .

IVUS analysis was performed according to the American College of Cardiology Clinical Expert Consensus Document on Standards for Acquisition, Measurement and Reporting of Intravascular Ultrasound Studies [6]. EEM and lumen CSAs were measured. Plaque plus media (P&M) CSA was calculated as EEM minus lumen CSA; and plaque burden was calculated as P&M divided by EEM CSA. Proximal and distal references were the single slices

with the largest lumen and smallest plaque CSAs within 10 mm proximally and distally, but before any large side branch. The lesion was the site with the smallest lumen CSA; if there were multiple image slices with the same minimum lumen CSA, then the image slice with the largest EEM and P&M was measured. Coronary artery remodeling was assessed by comparing the lesion site to the reference EEM CSA. Remodeling index was the lesion site EEM CSA divided by the average of the proximal and distal reference EEM CSA. PR was defined as a remodeling index >1.05 , intermediate remodeling (IR) as a remodeling index between 0.95 and 1.05, and NR as a remodeling index <0.95 [20]. Hypoechoic plaque was less bright compared with the reference adventitia. Hyperechoic, noncalcified plaque was as bright as or brighter than the reference adventitia without acoustic shadowing. Calcium plaque was hyperechoic with shadowing. A calcified lesion contained $>90^{\circ}$ of circumferential lesion calcium. When there was no dominant plaque composition, the plaque was classified as mixed.

A ruptured plaque contained a cavity that communicated with the lumen with an overlying residual fibrous cap fragment. A fragmented and loosely adherent plaque without a distinct cavity and without a fibrous cap fragment was not considered a plaque rupture. Rupture sites separated by a length of artery containing smooth lumen contours without cavities were considered to represent different plaque ruptures [21,22]. Plaque cavity was measured and extrapolated to the ruptured capsule area. Thrombus was an intraluminal mass having a layered or lobulated appearance, evidence of blood flow (microchannels) within the mass, and speckling or scintillation [22,23]. A lipid-pool like image was defined as a pooling of hypoechoic or echolucent material covered with a hyperechoic layer.

At post-intervention, we measured the minimum stent CSA. Percent stent expansion was calculated as minimum stent CSA divided by mean reference lumen CSA. Plaque prolapse was defined as tissue extrusion through the stent strut at post-intervention.

Statistical analysis

The statistical Package for Social Sciences (SPSS) for Windows, version 15.0 (Chicago, IL, USA) was used for all analyses. Continuous variables were presented as the mean value $\pm 1\text{SD}$; comparisons were conducted by Student's *t*-test or nonparametric Wilcoxon test if normality assumption was violated. Discrete variables were presented as percentages and relative frequencies; comparisons were conducted by chi-square statistics or Fisher's

Table 1 Baseline characteristics.

| | PR (n = 113) | IR/NR (n = 197) | p-value |
|---|--------------|-----------------|---------|
| Age (years) | 62 ± 13 | 66 ± 11 | 0.043 |
| Male gender | 72 (64) | 107 (54) | 0.11 |
| Clinical presentation | | | 0.9 |
| NSTEMI | 68 (61) | 117 (59) | |
| STEMI | 45 (40) | 80 (41) | |
| Diabetes mellitus | 46 (41) | 65 (33) | .17 |
| Hypertension | 89 (79) | 151 (77) | 0.7 |
| Smoking | 35 (31) | 70 (36) | 0.4 |
| Family history of coronary artery disease | 26 (23) | 26 (13) | 0.026 |
| Glycoprotein IIb/IIIa inhibitors | 26 (23) | 34 (17) | 0.2 |
| Use of distal protection devices | 13 (12) | 11 (6) | 0.060 |
| Ejection fraction (%) | 43 ± 13 | 46 ± 12 | 0.099 |
| cTnI (ng/ml) | 16.6 ± 47.4 | 14.4 ± 40.3 | 0.7 |
| White blood cells (10 ³ /mm ³) | 9.8 ± 4.1 | 8.8 ± 3.1 | 0.091 |
| Hemoglobin (g/dl) | 12.2 ± 2.2 | 12.3 ± 2.1 | 0.8 |
| Platelet count (10 ³ /mm ³) | 243 ± 89 | 222 ± 85 | 0.12 |
| Creatinine clearance (ml/min) | 60 ± 36 | 64 ± 31 | 0.3 |
| Total cholesterol (mg/dl) | 172 ± 47 | 168 ± 43 | 0.6 |
| Triglyceride (mg/dl) | 136 ± 49 | 125 ± 71 | 0.2 |
| LDL-cholesterol (mg/dl) | 105 ± 42 | 102 ± 37 | 0.6 |
| HDL-cholesterol (mg/dl) | 42 ± 12 | 43 ± 14 | 0.3 |
| High-sensitivity C-reactive protein (mg/dl) | 2.4 ± 5.4 | 1.9 ± 3.4 | 0.5 |

Data are n (%), or mean ± SD. PR, positive remodeling; IR/NR, intermediate remodeling/negative remodeling; NSTEMI, non-ST segment elevation myocardial infarction; STEMI, ST segment elevation myocardial infarction; cTnI, cardiac-specific troponin I; LDL, low-density lipoprotein; HDL, high-density lipoprotein.

exact test as appropriate. Multivariate analysis was performed to identify independent predictors of post-stenting cTnI elevation. Univariate analyses were first conducted to identify potential risk factors for post-stenting cTnI elevation. The likelihood ratio test was used, and the variables with a *p*-value of <0.2 were included in the multivariate model. Finally, a stepdown logistic regression was performed. The least significant variable was dropped at each step until only covariates with a *p*-value < 0.05 remained. A *p*-value < 0.05 was considered statistically significant.

Results

Patient characteristics

The baseline characteristics are summarized in Table 1. The PR group was younger and had more family history of coronary artery disease compared with the IR/NR group. There were trends toward more frequent use of distal protection devices, lower ejection fraction, and higher white blood cell counts in the PR group compared with the IR/NR group.

Angiographic and procedural results

Angiographic findings and procedural results are summarized in Table 2. Although there were no significant differences in infarct-related arteries, lesion location, number of diseased vessels, Thrombolysis In Myocardial Infarction (TIMI) flow grade, used stent types, and stent diameter and length between the PR group and the IR/NR group, angiographic lesion length was significantly longer and inflation pressure for stent implantation was significantly higher in the PR group compared with the IR/NR group. There was a strong trend toward increased PR when paclitaxel-eluting stents were used, compared with sirolimus-eluting or bare-metal stents [paclitaxel-eluting stent, 19/49 (39%); sirolimus-eluting stent, 35/138 (25%); bare-metal stent, 31/123 (25%); *p* = 0.15].

IVUS results

IVUS findings are summarized in Table 3. Lesion site EEM CSA, P&M CSA, and plaque burden were significantly greater in lesions with PR compared with lesions with IR/NR. IVUS lesion length was significantly longer, and calcium arc was signifi-

Table 2 Coronary angiographic findings and procedural results.

| | PR (<i>n</i> = 113) | IR/NR (<i>n</i> = 197) | <i>p</i> -value |
|---------------------------------|----------------------|-------------------------|-----------------|
| Infarct-related artery | | | 0.12 |
| Left main | 3 (3) | 2 (1) | |
| Left anterior descending artery | 54 (48) | 113 (57) | |
| Left circumflex artery | 13 (12) | 29 (15) | |
| Right coronary artery | 43 (38) | 53 (27) | |
| Lesion location | | | 0.096 |
| Ostium | 1 (1) | 3 (2) | |
| Proximal | 37 (33) | 84 (43) | |
| Middle | 67 (59) | 88 (45) | |
| Distal | 8 (7) | 22 (11) | |
| Multivessel disease | 54 (48) | 100 (51) | 0.4 |
| TIMI flow grade 0 | 23 (20) | 35 (18) | 0.5 |
| Stent type | | | 0.17 |
| Sirolimus-eluting stent | 53 (47) | 85 (43) | |
| Paclitaxel-eluting stent | 12 (11) | 37 (19) | |
| Bare-metal stent | 48 (43) | 75 (38) | |
| No. of deployed stents | 1.25 ± 0.53 | 1.16 ± 0.38 | 0.13 |
| Stent diameter (mm) | 3.31 ± 0.50 | 3.22 ± 0.41 | 0.10 |
| Stent length (mm) | 24.9 ± 11.6 | 23.6 ± 10.4 | 0.3 |
| Inflation pressure (mmHg) | 14.8 ± 2.7 | 14.2 ± 2.8 | 0.039 |
| Reference diameter (mm) | 3.26 ± 0.89 | 3.32 ± 0.75 | 0.5 |
| Pre-MLD (mm) | 0.63 ± 0.32 | 0.65 ± 0.51 | 0.7 |
| Lesion length (mm) | 21 ± 10 | 18 ± 8 | 0.031 |

Data are *n* (%), or mean ± SD. PR, positive remodeling; IR/NR, intermediate remodeling/negative remodeling; TIMI, thrombolysis in myocardial infarction; MLD, minimal lumen diameter.

Table 3 Intravascular ultrasound findings.

| | PR (<i>n</i> = 113) | IR/NR (<i>n</i> = 197) | <i>p</i> -value |
|--------------------------------------|----------------------|-------------------------|-----------------|
| Reference | | | |
| EEM CSA (mm ²) | 12.3 ± 4.5 | 13.2 ± 4.8 | 0.13 |
| Lumen CSA (mm ²) | 8.3 ± 3.2 | 9.0 ± 3.0 | 0.11 |
| P&M CSA (mm ²) | 4.0 ± 2.2 | 4.2 ± 2.8 | 0.6 |
| Plaque burden (%) | 32 ± 12 | 32 ± 11 | 1.0 |
| Lesion site | | | |
| EEM CSA (mm ²) | 14.6 ± 5.0 | 11.3 ± 3.9 | <0.001 |
| Lumen CSA (mm ²) | 2.6 ± 1.6 | 2.4 ± 1.0 | 0.7 |
| P&M CSA (mm ²) | 12.0 ± 4.5 | 8.9 ± 3.9 | <0.001 |
| Plaque burden (%) | 82 ± 36 | 76 ± 30 | 0.014 |
| IVUS lesion length (mm) | 26 ± 16 | 22 ± 12 | 0.026 |
| Calcium arc (°) | 97 ± 106 | 150 ± 106 | <0.001 |
| Lipid-pool like image | 45 (40) | 51 (26) | 0.011 |
| Scattered calcification | 70 (62) | 81 (41) | <0.001 |
| Plaque morphology | | | <0.001 |
| Hypoechoic | 72 (64) | 80 (41) | |
| Hyperechoic, non-calcified | 21 (19) | 31 (16) | |
| Hyperechoic, calcified | 12 (11) | 66 (34) | |
| Mixed | 8 (7) | 20 (10) | |
| Remodeling index | 1.20 ± 0.18 | 0.88 ± 0.13 | <0.001 |
| Minimum stent CSA (mm ²) | 7.92 ± 2.64 | 7.75 ± 2.56 | 0.079 |
| Stent expansion (%) | 95 ± 22 | 86 ± 29 | 0.034 |

Data are *n* (%), or mean ± SD. PR, positive remodeling; IR/NR, intermediate remodeling/negative remodeling; EEM, external elastic membrane; CSA, cross-sectional area; P&M, plaque plus media; IVUS, intravascular ultrasound.




Fig. 1 The incidence of plaque rupture and multiple plaque ruptures in lesions with positive remodeling (PR) vs. lesions with intermediate remodeling (IR)/negative remodeling (NR).

cantly smaller in lesions with PR compared with lesions with IR/NR. The presence of lipid-pool like image, scattered calcification, and hypoechoic plaque were significantly more common in lesions with PR compared with lesions with IR/NR. Percent stent expansion was significantly greater in lesions with PR compared with lesions with IR/NR.

Plaque rupture and multiple plaque ruptures were observed more frequently in lesions with PR compared with lesions with IR/NR (Fig. 1). Plaque cavity CSA was significantly greater and ruptured plaque length was significantly longer in lesions with PR compared with lesions with IR/NR (Fig. 2). IVUS-detected thrombus was observed more frequently in lesions with PR compared with lesions with IR/NR (Fig. 3). At post-intervention, plaque prolapse was observed more frequently in lesions with PR compared with lesions with IR/NR (Fig. 4).

Post-stenting cTnI elevation

Overall, cTnI was elevated after stenting in 179 patients (58% of the enrolled patients). cTnI was more significantly elevated after stenting in patients with PR compared with patients without IR/NR (Δ cTnI; $+7.8 \pm 51.1$ ng/ml vs. $+0.9 \pm 41.1$ ng/ml, $p=0.008$). Multiple logistic regression analysis was performed to determine independent predictors of post-stenting cTnI elevation. The following variables were tested (all with $p < 0.2$ in univariate analysis): male gender, diabetes mellitus, TIMI flow grade 0, stent length, IVUS plaque burden at the minimum lumen sites, plaque rupture, IVUS-detected thrombus, PR, plaque prolapse, and percent stent expansion. PR [odds ratio (OR) 1.92; 95% confidence interval




Fig. 2 Plaque cavity area (A) and ruptured plaque length (B) in lesions with positive remodeling (PR) vs. lesions with intermediate remodeling (IR)/negative remodeling (NR).

(CI) 1.04–2.98, $p=0.028$], plaque rupture (OR 1.98; 95% CI 1.16–3.45, $p=0.025$), IVUS-detected thrombus (OR 2.30; 95% CI 1.22–3.98, $p=0.008$), and plaque prolapse (OR 8.40; 95% CI 4.19–16.84, $p < 0.001$) were independently associated with post-stenting cTnI elevation.




Fig. 3 The incidence of intravascular ultrasound (IVUS)-detected thrombus in lesions with positive remodeling (PR) vs. lesions with intermediate remodeling (IR)/negative remodeling (NR).




Fig. 4 The incidence of post-stenting plaque prolapse in lesions with positive remodeling (PR) vs. lesions with intermediate remodeling (IR)/negative remodeling (NR).

Discussion

The present IVUS study demonstrated that AMI patients with PR had: 1) greater lesion site plaque burden, longer lesion length, and less severe calcification; 2) more lipid-pool like image, more scattered calcification, and more hypoechoic plaque; 3) greater percent stent expansion; 4) more plaque rupture and multiple plaque ruptures and more IVUS-detected thrombus; 5) more post-stenting plaque prolapsed; and 6) more significant cTnI elevation after stenting. Therefore, AMI patients with PR have more plaque vulnerability and higher frequency of plaque prolapse accompanied by more post-procedural myonecrosis compared with AMI patients with IR/NR.

Coronary arteries respond to plaque growth by either outward expansion of the vessel wall (PR) [1,2] or vessel shrinkage (NR) [3–5]. Several studies have suggested that lesions with PR may be both beneficial (avoiding luminal stenosis) and harmful (promoting plaque vulnerability) [8,24,25]. In contrast, lesions with NR may be associated with significant stenoses [5,26], but may appear more stable. Lesions with PR are also associated with higher long-term target lesion revascularization after nonstent interventional procedures [27], restenosis after bare-metal stent implantation [28–31], and drug-eluting stent implantation [32].

Autopsy studies have suggested that AMI results from spontaneous plaque rupture or erosion and subsequent thrombosis [33,34]. Several IVUS studies have reported varying percentage of culprit-lesion plaque rupture in patients with acute coronary syndrome [35–37]. Plaque ruptures tend to occur at a point where the fibrous cap is thinnest and most heavily infiltrated by macrophages indi-

cating ongoing inflammation at the site of plaque disruption [38]. Pathological studies have shown that coronary artery plaques with PR have a higher lipid content and macrophage count which are recognized histological markers for plaque vulnerability [39–41]. In the present study, lesions with PR have more plaque rupture, more multiple plaque ruptures, more IVUS-detected thrombus, more hypoechoic plaque, more lipid-pool like images, and more spotty calcification compared with lesions with IR/NR. The results of the present study are consistent with previous pathological and IVUS studies in which PR may make the plaque more vulnerable to rupture and cause AMI.

Plaque prolapse is an intraluminal tissue extrusion through stent struts and this can be easily and frequently detected by IVUS. It has been known that several pre-intervention IVUS features such as hypoechoic plaque rather than fibrous or calcified plaque, smaller minimal lumen diameter, and a larger plaque burden were associated with plaque prolapse, and the risk of plaque prolapse was higher during aggressive stenting procedure [35,36]. Our previous study showed that the incidence of plaque prolapse after stenting for infarct-related artery was 27% and pre-intervention IVUS lesion characteristics – plaque rupture and PR – and longer stent length predict plaque prolapse [42]. In the present study, plaque prolapse was observed more frequently in lesions with PR compared with lesions with IR/NR. Vulnerable plaques like PR and plaque rupture may provide the conditions for tissue extrusion through the stent struts.

Previous studies demonstrated that cardiac enzyme elevation after percutaneous coronary intervention is associated with higher mortality [43–47]. Cardiac enzyme elevation after coronary artery intervention is related to the large plaque burden and unstable plaque morphometry [48–50]. In addition to these pre-intervention IVUS findings, more aggressive stenting procedures appeared to be associated with greater levels of cardiac enzyme elevation [43,50–52]. In the present study, pre-intervention IVUS findings – PR, plaque rupture, IVUS-detected thrombus – and post-intervention plaque prolapse were independently associated with cTnI elevation after stent implantation. The results of the present study suggest that aggressive stenting with high pressure inflation for lesions with vulnerable plaque like PR, plaque rupture, and thrombus should be avoided to prevent post-stenting myonecrosis in patients with AMI because plaque prolapse is associated with aggressive stenting procedures and post-stenting cardiac enzyme elevation [35,42].

Study limitations

First, the present study is retrospective single-center study, so, is subject to limitations inherent in this type of clinical investigation. Second, IVUS imaging was performed at the discretion of the individual operators leading to potential selection bias. Third, it may be difficult to differentiate between an organized thrombus and plaque prolapse. IVUS data were analyzed by 2 independent observers (H.Y.J and C.Y.H). The presence of plaque prolapse was confirmed when 2 independent observers agreed. However, if so, it is possible that some cases identified as having plaque prolapse actually had thrombus prolapse. Fourth, volumetric IVUS analysis was not performed in the present study. Fifth, we excluded the patients with serious conditions such as severe heart failure, cardiogenic shock, and coronary artery bypass graft failure. Thus, the present study might not represent the whole spectrum of AMI patients. Sixth, long-term clinical follow-up was not available.

Conclusions

AMI patients with PR have more plaque vulnerability (more plaque rupture, more multiple plaque ruptures, and more IVUS-detected thrombus) and higher frequency of post-stenting plaque prolapse accompanied by more post-procedural cardiac enzyme elevation compared with AMI patients with IR/NR.

References

- [1] Glagov S, Weisenberg E, Zarins CK, Stankunavicius R, Koletis G. Compensatory enlargement of human atherosclerotic coronary arteries. *N Engl J Med* 1987;316:1371–5.
- [2] McPherson DD, Sirna SJ, Hiratzka LF, Thorpe L, Armstrong ML, Marcus ML, Kerber RE. Coronary arterial remodeling studied by high-frequency epicardial echocardiography: an early compensatory mechanism in patients with obstructive coronary atherosclerosis. *J Am Coll Cardiol* 1991;17:79–86.
- [3] Pasterkamp G, Wensing PJ, Post MJ, Hillen B, Mali WP, Borst C. Paradoxical arterial wall shrinkage may contribute to luminal narrowing of human atherosclerotic femoral arteries. *Circulation* 1995;91:1444–9.
- [4] Nishioka T, Luo H, Eigler NL, Berglund H, Kim CJ, Siegel RJ. Contribution of inadequate compensatory enlargement to development of human coronary artery stenosis: an in vivo intravascular ultrasound study. *J Am Coll Cardiol* 1996;27:1571–6.
- [5] Mintz GS, Kent KM, Pichard AD, Satler LF, Popma JJ, Leon MB. Contribution of inadequate arterial remodeling to the development of focal coronary artery stenoses. An intravascular ultrasound study. *Circulation* 1997;95:1791–8.
- [6] Mintz GS, Nissen SE, Anderson WD, Bailey SR, Erbel R, Fitzgerald PJ, Pinto FJ, Rosenfield K, Siegel RJ, Tuzcu EM, Yock PG. American College of Cardiology clinical expert consensus document on standards for acquisition, measurement and reporting of intravascular ultrasound studies (IVUS): a report of the American College of Cardiology task force on clinical expert consensus documents. *J Am Coll Cardiol* 2001;37:1478–92.
- [7] Birnbaum Y, Fishbein MC, Luo H, Nishioka T, Siegel RJ. Regional remodeling of atherosclerotic arteries: a major determinant of clinical manifestations of disease. *J Am Coll Cardiol* 1997;30:1149–64.
- [8] Schoenhagen P, Ziada KM, Kapadia SR, Crowe TD, Nissen SE, Tuzcu EM. Extent and direction of arterial remodeling in stable versus unstable coronary syndromes: an intravascular ultrasound study. *Circulation* 2000;101:598–603.
- [9] Fukuda D, Kawarabayashi T, Tanaka A, Nishibori Y, Taguchi H, Nishida Y, Shimada K, Yoshikawa J. Lesion characteristics of acute myocardial infarction: an investigation with intravascular ultrasound. *Heart* 2001;85:402–6.
- [10] Rioufol G, Finet G, Ginon I, André-Fouët X, Rossi R, Vialle E, Desjoyaux E, Convert G, Huret JF, Tabib A. Multiple atherosclerotic plaque rupture in acute coronary syndrome: a three-vessel intravascular ultrasound study. *Circulation* 2002;106:804–8.
- [11] Kotani J, Mintz GS, Castagna MT, Pinnow E, Berzinger CO, Bui AB, Pichard AD, Satler LF, Suddath WO, Waksman R, Laird Jr JR, Kent KM, Weissman NJ. Intravascular ultrasound analysis of infarct-related and non-infarct-related arteries in patients who presented with an acute myocardial infarction. *Circulation* 2003;107:2889–93.
- [12] Sano T, Tanaka A, Namba M, Nishibori Y, Nishida Y, Kawarabayashi T, Fukuda D, Shimada K, Yoshikawa J. C-reactive protein and lesion morphology in patients with acute myocardial infarction. *Circulation* 2003;108:282–5.
- [13] Hong MK, Mintz GS, Lee CW, Kim YH, Lee SW, Song JM, Han KH, Kang DH, Song JK, Kim JJ, Park SW, Park SJ. Comparison of coronary plaque rupture between stable angina and acute myocardial infarction: a three-vessel intravascular ultrasound study in 235 patients. *Circulation* 2004;110:928–33.
- [14] Tanaka A, Shimada K, Sano T, Namba M, Sakamoto T, Nishida Y, Kawarabayashi T, Fukuda D, Yoshikawa J. Multiple plaque rupture and C-reactive protein in acute myocardial infarction. *J Am Coll Cardiol* 2005;45:1594–9.
- [15] Hasegawa T, Ehara S, Kobayashi Y, Kataoka T, Yamashita H, Nishioka H, Asawa K, Yamagishi H, Yoshiyama M, Takeuchi K, Yoshikawa J, Ueda M. Acute myocardial infarction: clinical characteristics and plaque morphology between expansive remodeling and constrictive remodeling by intravascular ultrasound. *Am Heart J* 2006;151:332–7.
- [16] Ehara S, Kobayashi Y, Yoshiyama M, Shimada K, Shimada Y, Fukuda D, Nakamura Y, Yamashita H, Yamagishi H, Takeuchi K, Naruko T, Haze K, Becker AE, Yoshikawa J, Ueda M. Spotty calcification typifies the culprit plaque in patients with acute myocardial infarction: an intravascular ultrasound study. *Circulation* 2004;110:3424–9.
- [17] Virmani R, Kolodgie FD, Burke AP, Farb A, Schwartz SM. Lessons from sudden coronary death: a comprehensive morphological classification scheme for atherosclerotic lesions. *Arterioscler Thromb Vasc Biol* 2000;20:1262–75.
- [18] Falk E, Shah PK, Fuster V. Coronary plaque disruption. *Circulation* 1995;92:657–71.
- [19] Roberts WL, Moulton L, Law TC, Farrow G, Cooper-Anderson M, Savory J, Rifai N. Evaluation of nine automated high-sensitivity C-reactive protein methods: implications for

- clinical and epidemiological applications. Part 2. *Clin Chem* 2001;47:418–25.
- [20] Nakamura M, Nishikawa H, Mukai S, Setsuda M, Nakajima K, Tamada H, Suzuki H, Ohnishi T, Kakuta Y, Nakano T, Yeung AC. Impact of coronary artery remodeling on clinical presentation of coronary artery disease: an intravascular ultrasound study. *J Am Coll Cardiol* 2001;37:63–9.
 - [21] Maehara A, Mintz GS, Bui AB, Walter OR, Castagna MT, Canos D, Pichard AD, Satler LF, Waksman R, Suddath WO, Laird Jr JR, Kent KM, Weissman NJ. Morphologic and angiographic features of coronary plaque rupture detected by intravascular ultrasound. *J Am Coll Cardiol* 2002;40:904–10.
 - [22] Fujii K, Kobayashi Y, Mintz GS, Takebayashi H, Dangas G, Moussa I, Mehran R, Lansky AJ, Kreps E, Collins M, Colombo A, Stone GW, Leon MB, Moses JW. Intravascular ultrasound assessment of ulcerated ruptured plaques: a comparison of culprit and nonculprit lesions of patients with acute coronary syndromes and lesions in patients without acute coronary syndromes. *Circulation* 2003;108:2473–8.
 - [23] Chemarin-Alibelli MJ, Pieraggi MT, Elbaz M, Carrié D, Fourcade J, Puel J, Tobis J. Identification of coronary thrombus after myocardial infarction by intracoronary ultrasound compared with histology of tissues sampled by atherectomy. *Am J Cardiol* 1996;77:344–9.
 - [24] Smits PC, Pasterkamp G, de Jaegere PP, de Feyter PJ, Borst C. Angioscopic complex lesions are predominantly compensatory enlarged: an angiography and intracoronary ultrasound study. *Cardiovasc Res* 1999;41:458–64.
 - [25] Pasterkamp G, Schoneveld AH, van der Wal AC, Haudenschild CC, Clarijs RJ, Becker AE, Hillen B, Borst C. Relation of arterial geometry to luminal narrowing and histologic markers for plaque vulnerability: the remodeling paradox. *J Am Coll Cardiol* 1998;32:655–62.
 - [26] Pasterkamp G, Wensing PJ, Hillen B, Post MJ, Mali WP, Borst C. Impact of local atherosclerotic remodeling on the calculation of percent luminal narrowing. *Am J Cardiol* 1997;79:402–5.
 - [27] Dangas G, Mintz GS, Mehran R, Lansky AJ, Kornowski R, Pichard AD, Satler LF, Kent KM, Stone GW, Leon MB. Preintervention arterial remodeling as an independent predictor of target-lesion revascularization after nonstent coronary intervention: an analysis of 777 lesions with intravascular ultrasound imaging. *Circulation* 1999;99:3149–54.
 - [28] Endo A, Hirayama H, Yoshida O, Arakawa T, Akima T, Yamada T, Nanasato M. Arterial remodeling influences the development of intimal hyperplasia after stent implantation. *J Am Coll Cardiol* 2001;37:70–5.
 - [29] Okura H, Morino Y, Oshima A, Hayase M, Ward MR, Popma JJ, Kuntz RE, Bonneau HN, Yock PG, Fitzgerald PJ. Preintervention arterial remodeling affects clinical outcome following stenting: an intravascular ultrasound study. *J Am Coll Cardiol* 2001;37:1031–5.
 - [30] Sahara M, Kirigaya H, Oikawa Y, Yajima J, Ogasawara K, Satoh H, Nagashima K, Hara H, Nakatsu Y, Aizawa T. Arterial remodeling patterns before intervention predict diffuse in-stent restenosis: an intravascular ultrasound study. *J Am Coll Cardiol* 2003;42:1731–8.
 - [31] Hong YJ, Jeong MH, Hyun DW, Hur SH, Kim KB, Kim W, Lim SY, Lee SH, Hong SN, Kim KH, Yun KH, Kang DG, Lee YS, Park HW, Kim JH, et al. Impact of preinterventional arterial remodeling on in-stent neointimal hyperplasia and in-stent restenosis after coronary stent implantation. *Circ J* 2005;69:414–9.
 - [32] Mintz GS, Tinana A, Hong MK, Lee CW, Kim JJ, Farnot NE, Park SW, Park SJ, Weissman NJ. Impact of preinterventional arterial remodeling on neointimal hyperplasia after implantation of (non-polymer-encapsulated) paclitaxel-coated stents: a serial volumetric intravascular ultrasound analysis from the ASian Paclitaxel-Eluting Stent Clinical Trial (ASPECT). *Circulation* 2003;108:1295–8.
 - [33] Ponde CK, Aroney CN, McEniery PT, Bett JH. Plaque prolapse between the struts of the intracoronary Palmaz-Schatz stent: report of two cases with a novel treatment of this unusual problem. *Cathet Cardiovasc Diagn* 1997;40:353–7.
 - [34] Brack MJ, Forbat LN, Skehan JD, Gershlick AH. Plaque herniation through an intracoronary stent. *Int J Cardiol* 1994;44:93–5.
 - [35] Bocksch W, Scharf M, Beckmann S, Dreyse S, Fleck E. Intravascular ultrasound imaging in patients with acute myocardial infarction. *Eur Heart J* 1995;16(Suppl.):46–52.
 - [36] Hong MK, Park SW, Lee CW, Kang DH, Song JK, Kim JJ, Park SJ. Long-term outcomes of minor plaque prolapsed within stents documented with intravascular ultrasound. *Catheter Cardiovasc Interv* 2000;51:22–6.
 - [37] Cheneau E, Leborgne L, Mintz GS, Kotani J, Pichard AD, Satler LF, Canos D, Castagna M, Weissman NJ, Waksman R. Predictors of subacute stent thrombosis: results of a systematic intravascular ultrasound study. *Circulation* 2003;108:43–7.
 - [38] Farb A, Burke AP, Kolodgie FD, Virmani R. Pathological mechanisms of fatal late coronary stent thrombosis in humans. *Circulation* 2003;108:1701–6.
 - [39] Varnava AM, Mills PG, Davies MJ. Relationship between coronary artery remodeling and plaque vulnerability. *Circulation* 2002;105:939–43.
 - [40] Davies MJ, Richardson PD, Woolf N, Katz DR, Mann J. Risk of thrombosis in human atherosclerotic plaques: role of extracellular lipid, macrophage, and smooth muscle cell content. *Br Heart J* 1993;69:377–81.
 - [41] Moreno PR, Falk E, Palacios IF, Newell JB, Fuster V, Fallon JT. Macrophage infiltration in acute coronary syndromes: implications for plaque rupture. *Circulation* 1994;90:775–8.
 - [42] Hong YJ, Jeong MH, Ahn Y, Sim DS, Chung JW, Cho JS, Yoon NS, Yoon HJ, Moon JY, Kim KH, Park HW, Kim JH, Cho JG, Park JC, Kang JC. Incidence, predictors, and outcome of plaque prolapse after stent implantation in patients with acute myocardial infarction: an intravascular ultrasound analysis. *J Am Coll Cardiol* 2008;1:489–97.
 - [43] Harrington RA, Lincoff AM, Califf RM, Holmes Jr DR, Berdan LG, O'Hanesian MA, Keeler GP, Garratt KN, Ohman EM, Mark DB, Jacobs AK, Topol EJ. Characteristics and consequences of myocardial infarction after percutaneous coronary intervention: insights from the Coronary Angioplasty Versus Excisional Atherectomy Trial (CAVEAT). *J Am Coll Cardiol* 1995;25:1693–9.
 - [44] Tardiff BE, Califf RM, Tcheng JE, Lincoff AM, Sigmon KN, Harrington RA, Mahaffey KW, Ohman EM, Teirstein PS, Blankenship JC, Kitt MM, Topol EJ. Clinical outcomes after detection of elevated cardiac enzymes in patients undergoing percutaneous intervention. IMPACT-II investigators. integrilin (eptifibatide) to minimize platelet aggregation and coronary Thrombolysis-II. *J Am Coll Cardiol* 1999;33:88–96.
 - [45] Kong TQ, Davidson CJ, Meyers SN, Tauke JT, Parker MA, Bonow RO. Prognostic implication of creatine kinase elevation following elective coronary artery interventions. *JAMA* 1997;277:461–6.
 - [46] Kini A, Marmur JD, Kini S, Dangas G, Cocke TP, Wallenstein S, Brown E, Ambrose JA, Sharma SK. Creatine kinase-MB elevation after coronary intervention correlates with diffuse atherosclerosis, and low-to-medium level elevation has a benign clinical course: implications for early discharge after coronary intervention. *J Am Coll Cardiol* 1999;34:663–71.

- [47] Saucedo JF, Mehran R, Dangas G, Hong MK, Lansky A, Kent KM, Satler LF, Pichard AD, Stone GW, Leon MB. Long-term clinical events following creatine kinase—myocardial band isoenzyme elevation after successful coronary stenting. *J Am Coll Cardiol* 2000;35:1134–41.
- [48] Furuichi S, Itoh A, Ishibashi-Ueda H, Noguchi T, Miyawaki M, Nomura K, Arita Y, Otsuka M, Shindo T. Ultrasound attenuated coronary plaque as a risk factor for slow flow or no-reflow during percutaneous coronary intervention: a case report. *J Cardiol* 2007;49:193–7.
- [49] Fujii K, Carlier SG, Mintz GS, Kobayashi Y, Jacoboff D, Nierenberg H, Takebayashi H, Yasuda T, Moussa I, Dangas G, Mehran R, Lansky AJ, Kreps EM, Collins M, Stone GW, et al. Creatine kinase-MB enzyme elevation and long-term clinical events after successful coronary stenting in lesions with ruptured plaque. *Am J Cardiol* 2005;95:355–9.
- [50] Mehran R, Dangas G, Mintz GS, Lansky AJ, Pichard AD, Satler LF, Kent KM, Stone GW, Leon MB. Atherosclerotic plaque burden and CK-MB enzyme elevation after coronary interventions: intravascular ultrasound study of 2256 patients. *Circulation* 2000;101:604–10.
- [51] Kugelmass AD, Cohen DJ, Moscucci M, Piana RN, Senerchia C, Kuntz RE, Baim DS. Elevation of the creatine kinase myocardial isoform following otherwise successful directional coronary atherectomy and stenting. *Am J Cardiol* 1994;74:748–54.
- [52] Califf RM, Abdelmeguid AE, Kuntz RE, Popma JJ, Davidson CJ, Cohen EA, Kleiman NS, Mahaffey KW, Topol EJ, Pepine CJ, Lipicky RJ, Granger CB, Harrington RA, Tardiff BE, Crenshaw BS, et al. Myonecrosis after revascularization procedures. *J Am Coll Cardiol* 1998;31:241–51.

Available online at www.sciencedirect.com



ScienceDirect